

## ROUTINE EYE DROPS for CATARACT SURGERY

Dr. Ross Littlewood

9374 0620

litwood@westnet.com.au

	<b>BREAKFAST</b>	<b>LUNCH</b>	<b>DINNER</b>	<b>BED TIME</b>
1 day <b>PRE OP</b>	Tobrex Acular	Tobrex Acular	Tobrex Acular	Tobrex Acular
<b>Day of surgery</b>	Tobrex Acular	Surgery	Tobrex Acular Maxidex	Tobrex Acular Maxidex
1 day <b>POST OP</b>	Tobrex Acular Maxidex	Tobrex Acular Maxidex	Tobrex Acular Maxidex	Tobrex Acular Maxidex
2 days POSTOP	Tobrex Acular Maxidex	Tobrex Acular Maxidex	Tobrex Acular Maxidex	Tobrex Acular Maxidex
3 days POSTOP	Tobrex Acular Maxidex	Tobrex Acular Maxidex	Tobrex Acular Maxidex	Tobrex Acular Maxidex
4 days POSTOP	Tobrex Acular Maxidex	Tobrex Acular Maxidex	Tobrex Acular Maxidex	Tobrex Acular Maxidex
5 days POSTOP	Maxidex	Maxidex	Maxidex	Maxidex
6 days POSTOP	Maxidex	Maxidex	Maxidex	Maxidex
7 days POSTOP	Maxidex			Maxidex
8 days POSTOP	Maxidex			Maxidex
9 days POSTOP	Maxidex			Maxidex
10 days POSTOP	Maxidex			Maxidex
11 days POSTOP	Maxidex			Maxidex
12 days POSTOP	Maxidex			Maxidex
13 days POSTOP	Maxidex			Maxidex
14 days POSTOP	Maxidex			Maxidex
Beyond 14 days	Until the bottle is empty			Maxidex
<b>If the eye is uncomfortable use preservative free lubricants frequently and contact the office for a further appointment.</b>				

**For surface irritation at any time use preservative free lubricants such as Optive Advanced, Optive Sensitive, Systane, Bion, or Hylo drops every 2 hours.**

**Eyes with pre-existing disease often require a modification to this routine schedule.**